

ASYMPTOTIC DIMENSION OF FINITELY PRESENTED GROUPS

THANOS GENTIMIS

ABSTRACT. We prove that if a finitely presented group is one-ended then its asymptotic dimension is bigger than 1. It follows that a finitely presented group of asymptotic dimension 1 is virtually free.

1. INTRODUCTION

The notion of asymptotic dimension of a metric space was introduced by Gromov in [6]. It is a large scale analog of topological dimension and it is invariant by quasi-isometries. This notion has proved relevant in the context of Novikov's higher signature conjecture and it was investigated further by other people (see [13], [1], [10]).

In this paper we show the following:

Theorem 1. *If G is a one-ended finitely presented group then G has asymptotic dimension greater or equal to 2.*

Also we deduce as corollary:

Theorem 2. *If G is a finitely presented group with $\text{asdim } G = 1$ then G is virtually free.*

For finitely generated groups the statement above doesn't hold. We give a counter-example at the end.

After we completed this work T.Januszkiewicz brought to our attention his joint paper with J.Swiatkowski ([8]) where they proved the same results independently.

Finally I would like to thank pr. P.Papasoglu for his help and guidance during the writing of this paper.

2. PRELIMINARIES

Metric Spaces. Let (X, d) be a metric space. If A, B are subsets of X we set $d(A, B) = \inf\{d(a, b) : a \in A, b \in B\}$. A *path* in X is a map $\gamma : I \rightarrow X$ where I is an interval in $\mathbb{R}$. A path γ joins two points x and y in X if $I = [a, b]$, $\gamma(a) = x$ and $\gamma(b) = y$. The path γ is called an *infinite ray* starting from x_0 if $I = [0, \infty)$ and $\gamma(0) = x_0$. A *geodesic*, a *geodesic ray*, or a *geodesic segment* in X , is an isometry $\gamma : I \rightarrow X$ where I is $\mathbb{R}$ or $[0, \infty)$ or a closed interval in $\mathbb{R}$. We use the terms geodesic, geodesic ray etc. for the images of γ without discrimination. On a path-connected space X given two points x, y we define the *path metric* to

Date: February 2, 2008.

Key words and phrases. dimension theory, asymptotic dimension, finitely presented group.

Research co-funded by the European Social Fund and National Resources-EPEAEK II-PYTHAGORAS.

be $\rho(x, y) = \inf\{\text{length}(p)\}$ where the infimum is taken over all paths p that join x and y . A space is called a *geodesic metric space* if for every x, y in X there exists a geodesic segment which joins them. In a geodesic space the path metric is indeed a metric. A geodesic metric space X is said to be *one-ended* if for every bounded K , $X - K$ has exactly one unbounded connected component. We say that X is *uniformly one-ended*, if for every $n \in \mathbb{R}^+$ there is an $m \in \mathbb{R}^+$ such that for every $K \subset X$ with $\text{diam}(K) < n$, $X - K$ has exactly one connected component of diameter bigger than m .

Groups. The *Cayley graph* of G with respect to a generating set S is the 1-dimensional complex having a vertex for each element of G and an edge joining vertex x to vertex xs for every vertex x and every $s \in S$. The Cayley graph has a natural metric which makes it a geodesic metric space, where each edge has length 1 (see [10]). In fact any connected graph can be made geodesic metric space in the same way.

We will use the same letter G for both the group and its Cayley graph as a metric space. We also use *van - Kampen Diagrams* (see [9] chapter V pg 236 -240 and [5]). A van Kampen diagram $\mathfrak{D}$ for a word w in S representing the identity element of G , is a finite, planar, contractible, combinatorial 2-complex; its 1-cells are directed and labeled by generators and the boundary labels of each of its 2-cells are cyclic conjugates of relators or inverse relators. Further the boundary label for $\mathfrak{D}$ is w when read (by convention anticlockwise) from a base point in $\partial\mathfrak{D}$. We recall here that a word w represents the identity element of G if and only if the path in the Cayley graph labeled by w is closed.

There is a natural map f from the 1-skeleton $\mathfrak{D}^{(1)}$ of $\mathfrak{D}$ to the Cayley graph of G . f sends the base point to a vertex v of the Cayley graph and edges of $\mathfrak{D}^{(1)}$ to edges of the Cayley graph with the same label. Obviously f is determined by the image of the base point, v . f is not necessarily injective. If we consider $\mathfrak{D}^{(1)}$ as a geodesic metric space giving each edge length 1 then $d(x, y) \geq d(f(x), f(y))$ for every x, y in $\mathfrak{D}^{(1)}$.

We say that a group G is *virtually free* if there exists a finite index subgroup H of G which is a free group.

A group G is called *free-by-finite* if it contains a normal subgroup N of finite index such that N is free.

Generally if X is a property of groups then we say that G is *virtually X* if $\exists H <_f G$ and H has property X . We say that G is *X -by-finite* if $\exists N <_f G$ normal such that N has property X .

Obviously if G is X -by-finite then G is *virtually X* . The converse also holds if the property X is inherited to subgroups.

Now since if G is free every subgroup of G is free we have that G is *virtually free* if and only if G is *free by-finite*.

Asymptotic Dimension. A metric space Y is said to be d - disconnected or that it has dimension 0 on the d - scale if there exist $B_i \subset Y$ such that:

$$Y = \bigcup_{i \in I} B_i$$

with $\sup\{\text{diam} B_i, i \in I\} \leq D < \infty, d(B_i, B_j) \geq d \forall i \neq j$.

Definition. (*Asymptotic Dimension 1*) We say that a space X has asymptotic dimension n if n is the minimal number such that for every $d > 0$ we have: $X = \bigcup X_k$ for $k = 0, 1, 2, \dots, n$ and all X_k are d -disconnected. We then write $\text{asdim } X = n$.

We say that a covering $\{B_i\}$ of X has d - multiplicity $\leq k$, if and only if every d -ball $B(x, d)$ in X meets no more than k sets B_i of the covering. The covering has d - multiplicity k if d - multiplicity $\leq k$ is true and d - multiplicity $\leq k - 1$ is false. A covering $\{B_i\}$, $i \in I$ is D - bounded, if $\text{diam}(B_i) \leq D$ for all $i \in I$.

Definition. (*Asymptotic Dimension 2*) We say that a space X has $\text{asdim } X = n$, if n is the minimal number such that $\forall d > 0$ there exists a D -bounded covering of X with d - multiplicity $\leq n + 1$.

The two definitions used here are the first two definitions Gromov gave in his paper [6]. It is not difficult to see that the two definitions are equivalent.

3. MAIN THEOREM

Before we get to the main theorem we will prove two lemmas that we will need below.

Lemma 1. *Let G be a finitely generated, infinite group then $\text{asdim } G > 0$.*

Proof. Let $\text{asdim } G = 0$ and fix $d > 0$. Then according to the first definition we have that $G = X_1$ where $X_1 = \bigcup B_i$ with:

- (1) $\text{diam } B_i \leq D, \forall i \in I$
- (2) $d(B_i, B_j) \geq d, \forall i \neq j$.

That means that G is d -disconnected. Since G is a connected graph it follows that we can not have two distinct B_i 's. So $G \subset B_1$ which means that G is D -bounded. But since G is finitely generated we have immediately that G is finite which is a contradiction. $\square$

Lemma 2. *If G is an one-ended finitely presented group then G contains a bi-infinite geodesic.*

Proof. Take any $n \in \mathbb{N}$. Then define

$$C_n = \{\text{geodesic paths starting from } e \text{ of length } n\}$$

We note that:

- a) $C_n \neq \emptyset$ for all $n \in \mathbb{N}$.
- b) C_n is a finite set for all $n \in \mathbb{N}$.

Consider a map

$$\pi_{n-1}^n : C_n \rightarrow C_{n-1}$$

which takes every geodesic path of length n from C_n and cuts off the last edge of that path. Then obviously what is left is also a geodesic and now the length is $n - 1$. Thus it is contained in C_{n-1} . So for every $n \in \mathbb{N}$, π_{n-1}^n is a natural well defined map.

So now consider the inverse limit sequence

$$\{e\} \leftarrow C_1 \leftarrow C_2 \leftarrow \dots \leftarrow C_n \leftarrow C_{n+1} \leftarrow \dots$$

with bonding maps π_{n-1}^n . By using the well known statement that the inverse limit of compact spaces is compact and hence nonempty, we obtain that

$$F = \varprojlim \{C_i, \pi_{i-1}^i\} \neq \emptyset$$

Now let $r \in F$. Obviously r is an infinite geodesic ray starting from e . Now let's fix a geodesic segment ξ in C_n and denote by Y_ξ all the infinite geodesics in F starting with ξ . Then obviously Y_ξ is the inverse image $(\pi_n^\infty)^{-1}(\xi)$ where $\pi_n^\infty : F \rightarrow C_n$ is the natural projection. Thus if Y_ξ is non empty it is compact.

Now define:

$$\begin{aligned} S_n &= \{g \in G \text{ such that } d(g, e) = n\}, \\ L_n &= \{\text{all the geodesic paths starting from } S_n \text{ and ending at } e\}, \\ E_n &= \{r : r \text{ are geodesic rays, } r(0) \in K_n, e \in r_i\} \end{aligned}$$

Obviously E_n is not empty since let r' one of the geodesic rays we defined starting from e . Let $g = r'(n)$ be the vertex of r' in S_n . Define $g^{-1} \cdot r'$. Since multiplying with an element doesn't change the respective distances $g^{-1} \cdot r'$ is also a geodesic starting from $g^{-1} \cdot e = g^{-1}$ which is obviously in S_n and passing through $g^{-1} \cdot g = e$. Thus, $E_n \neq \emptyset$.

Now fix an η in L_n and its corresponding $h \in S_n$. Define the inverse limit

$$\{h\} \leftarrow h \cdot C_1 \leftarrow h \cdot C_2 \leftarrow \dots \leftarrow h \cdot C_n \leftarrow \dots$$

The above inverse limit gives us all the geodesic rays starting from h i.e. $h \cdot F$.

The set Y'_η of all geodesic rays starting with the path η being homeomorphic to $Y_{h^{-1} \cdot \eta}$, is a compact by the above remark (if it is not empty of course!).

Then by the definition we get that:

$$E_n = \bigcup_{\eta \in L_n} Y'_\eta$$

Since E_n is non-empty, it is compact as a finite union of compact sets. Thus the inverse limit:

$$E = \varprojlim E_n$$

with the restrictions as the bonding maps is compact and nonempty. Now let k in E . Then k is a path that passes through e and k is our bi-infinite geodesic line. This concludes the lemma. $\square$

Theorem 1. *If G is an one-ended finitely presented group then $\text{asdim } G \geq 2$.*

Proof. By lemma 1 and since G is one-ended we have that G is infinite and thus $\text{asdim } G > 0$. We will show that $\text{asdim } G \neq 1$. Let's suppose that $\text{asdim } G = 1$.

Let $M = \max\{|r_i|, i = 1, 2, \dots, n: r_i \text{ relation of } G\}$, where $|r| = \text{length}$ of the word r . We fix $d > 100M + 100$. Since $\text{asdim } G = 1$ there is a covering $\mathbb{B} = \{B_i\}$ with:

$$G = \bigcup_{i \in I} B_i$$

and $\text{diam } B_i < D, \forall i \in I$, such that every ball $B(x, d)$ intersects at most 2 sets of the covering $\mathbb{B}$. We may assume without loss of generality that if r is a path in the Cayley graph labeled by a relator r_i then r is contained in some $B_j \in \mathbb{B}$.

Since G is one-ended we have that G has a bi-infinite geodesic S (Lemma 2). Let $N = \max\{100D^{100}, 300M\}$. Choose an $x_0 \in S$ and consider the ball $B(x_0, N)$

which separates the geodesic S into two geodesic rays S_1 and S_2 . Since G is one-ended there is an x in S_1 , a y in S_2 and a path p with $p(0) = x$ and $p(t) = y$ such that $p \cap B(x_0, N) = \emptyset$.

We denote by $[x, y]$ the part of the geodesic S , that connects x and y . Obviously $\text{length}([x, y]) \geq 2N$. We denote by w the path that corresponds to $[x, y] \cup p$. We have then that

$$\text{length}(w) = \text{length}([x, y]) + \text{length}(p) \Rightarrow \text{length}(w) > 200D$$

So in order to cover the path w we need at least 3 sets of the covering $\{B_i\}$.

We consider now the van-Kampen diagram $\mathfrak{D}$ that corresponds to the path w and the function f from $\mathfrak{D}^{(1)}$ to the Cayley graph G . So $f(\partial\mathfrak{D}) = w$. For notational convenience we label vertices and edges of $\partial\mathfrak{D}$ in the same way as w . So for example we denote the vertex on $\partial\mathfrak{D}$ which is mapped to $x_0 \in w$ by f also by x_0 .

Let B be a set of the covering that intersects $[x, y]$. We consider $f^{-1}(B)$. Let $C(B)$ be the union of all 2-cells of $\mathfrak{D}$ which have the property that their boundary is contained in $f^{-1}(B)$. Let U be the collection of all such sets B with the following property: For some connected component, K , of $C(B)$, $[x, y] \cap K$ is contained in an interval $[a, b]$ with $a, b \in K$ such that $x_0 \in [a, b]$. Let $d(K) = d(a, b)$ for such a component and let $d(B)$ be the maximal value of all $d(K)$ for K component of $C(B)$ such that $x_0 \in [a, b]$. Let B_1 be a set in U for which $d(B_1)$ is maximal. Let K_1 be the connected component of $C(B_1)$ for which $d(K_1) = d(B_1)$. Let's say that $K_1 \cap [x, y]$ is contained in $[a_1, b_1]$ with $a_1, b_1 \in K_1$. Let e be the edge of $[x, y]$ adjacent to a_1 which does not lie in $[a, b]$. If r is the 2-cell containing e there is some $B_2 \in \mathbb{B}$ such that $C(B_2)$ contains r .

Let C be the subset of $\mathfrak{D}$ which contains all 2-cells with boundary contained in $f^{-1}(B_1 \cup B_2)$. Since $d(x_0, p) \geq N$, C does not intersect p . Thus $\mathfrak{D} - C \neq \emptyset$. Let K be the connected component of C which contains x_0 .

Let $P = \mathfrak{D} - K \cap \overline{K}$. P is connected since K is connected. Each edge of P is contained in two 2-cells. One of these 2-cells lies in $f^{-1}(B_1 \cup B_2)$ and one does not lie in this set. It is not possible that all edges of P are contained in a 2-cell of $f^{-1}(B_2)$. Indeed in this case we would have $d(B_2) > d(B_1)$, which is impossible. Since r is contained in $C(B_2)$ some edge of P is not contained in a 2-cell of $f^{-1}(B_1)$. It follows that there are 2 adjacent edges e_1, e_2 in P such that one of them is contained in a 2-cell of $f^{-1}(B_1)$ and the other in a 2-cell of $f^{-1}(B_2)$. If c is the 2-cell that contains e_1 and is not in $f^{-1}(B_1 \cup B_2)$ then c lies in a set $f^{-1}(B_3)$ with $B_3 \in \mathbb{B}$ and $B_3 \neq B_1, B_2$. The edges e_1, e_2 and the 2-cell c have a vertex v in common. So $v \in f^{-1}(B_1 \cap B_2 \cap B_3)$. It follows that $B_1 \cap B_2 \cap B_3 \neq \emptyset$, a contradiction.

This concludes the proof. $\square$

Remark. *The result above holds also for uniformly one-ended simply connected simplicial complexes. So if X is a uniformly one-ended simply connected simplicial complex then $\text{asdim } X \geq 2$.*

Of course the result does not hold for one-ended simply connected simplicial complexes, a half-line gives a counterexample. We remark finally that if a Cayley graph is one ended then it is uniformly one ended.

We note that the following theorem holds:

Theorem. (*Dunwoody-Stallings [4]*) *If G is a finitely presented group then G is the fundamental group of a graph of groups such that all the edge groups are finite and all the vertex groups are 0 or 1 ended.*

Also it is known that:

Lemma 3. *If all the vertex groups are 0-ended (i.e. finite) then G is virtually free (see [11], page 120, prop.11).*

Furthermore it is not difficult to prove the following lemma (see [7]):

Lemma 4. *If $H < G$ and H is finitely generated then $\text{asdim } G \geq \text{asdim } H$.*

Using the lemma above and theorem 1 we have the stronger result:

Theorem 2. *If G is a finitely presented group with $\text{asdim } G = 1$ then G is virtually free.*

Proof. Let G be a finitely presented group with $\text{asdim } G = 1$. Let Γ be the graph of groups of the Dunwoody-Stallings theorem. If a vertex group H is one-ended then from the theorem 1, $\text{asdim } H \geq 2$. But $H < G$ which means that $\text{asdim } G \geq 2$ which is a contradiction. So all vertex groups are 0-ended. It follows that G is virtually free. $\square$

Now we give an example of a finitely generated group which is not finitely presented, not virtually free and that has asymptotic dimension 1. Namely:

Proposition. *Let $G = \mathbb{Z}_2 \wr_r \mathbb{Z}$ be the restricted wreath product of $\mathbb{Z}_2$ and $\mathbb{Z}$. Then $\text{asdim } G = 1$ and G is not virtually free.*

Proof. Since $G = \mathbb{Z}_2 \wr_r \mathbb{Z}$ there exists a short exact sequence:

$$0 \rightarrow (\oplus_{\mathbb{Z}} \mathbb{Z}_2) \rightarrow G \rightarrow \mathbb{Z} \rightarrow 0$$

By the Hurewicz type formula (see [3]) we have:

$$\text{asdim } G \leq \text{asdim } \mathbb{Z} + \text{asdim}(\oplus_{\mathbb{Z}} \mathbb{Z}_2)$$

Since every finitely generated subgroup F of $(\oplus_{\mathbb{Z}} \mathbb{Z}_2)$ is finite, we have that all F of that type have asymptotic dimension 0. Following the definition of asymptotic dimension for arbitrary discrete groups found in [3] we get:

$$\text{asdim}(\oplus_{\mathbb{Z}} \mathbb{Z}_2) = \sup\{\text{asdim } F \mid F < G, \text{ finitely generated}\} = 0$$

Another way to get the same result is by using the following corollary found in [12]

Corollary. *Let G be a countable abelian group. Then $\text{asdim } G = 0$ if and only if G is torsion.*

Obviously $\oplus_{\mathbb{Z}} \mathbb{Z}_2$ is abelian and torsion so $\text{asdim } \oplus_{\mathbb{Z}} \mathbb{Z}_2 = 0$. Thus we get:

$$\text{asdim } G \leq 1 + 0 = 1$$

Since G is a finitely generated infinite group (see [2] for the description of the generators), by lemma 1 we have that $\text{asdim } G > 0$. So $\text{asdim } G = 1$.

We will prove that G is not virtually free. Let G be virtually free. Then G is free-by-finite. So there exists a normal subgroup N of G such that N is free and the index $|G : N|$ is finite. Recall the exact sequence:

$$0 \rightarrow (\oplus_{\mathbb{Z}} \mathbb{Z}_2) \rightarrow G \rightarrow \mathbb{Z} \rightarrow 0$$

and let's denote the image of $\oplus \mathbb{Z} \mathbb{Z}_2$ in G to be H through the one to one mapping f . Then obviously $H < G$. Define $q : G \rightarrow \frac{G}{N}$ and restrict q to H . Since H is infinite and $\frac{G}{N}$ is finite then $H \cap \ker q \neq \emptyset$. But $\ker q = N$ thus $H \cap N \neq \emptyset$ which is a contradiction because if $a \in H$ a is torsion. Thus $a \notin N$ since N is free. $\square$

REFERENCES

1. A.Dranishnikov, *Asymptotic topology*, Russian Math.Surveys 55(2000), No 6, 71-116.
2. A.Dranishnikov, *Groups With Polynomial Dimension Growth*, Geometriae Dedicata, arxiv: math.MG/0405239
3. A.Dranishnikov and J.Smith, *Asymptotic Dimension of Discrete Groups*, Fundamenta Mathematicae, 189 (2006)
4. M.Dunwoody, *The accessibility of finitely presented groups*, Invent. Math. 81 (1985), No 3, 449-457.
5. S.Gersten and Tim Riley, *Filling length in Finitely presentable groups*, Geometriae Dedicata 92: 4158, 2002.
6. M.Gromov, *Asymptotic invariants of infinite groups*, 'Geometric group theory', (G.Niblo, M.Roller, Eds.), LMS Lecture Notes, vol. 182, Cambridge Univ. Press (1993)
7. M.Gromov, *Hyperbolic groups*, Essays in group theory (S. M. Gersten, ed.), MSRI Publ. 8, Springer-Verlag, 1987 pp. 75-263.
8. T.Januszkiewicz and J.Swiatkowski, *Filling Invariants In Systolic Complexes and Groups* "preprint" (June 2005).
9. R.Lyndon, P.Schupp, *Combinatorial Group Theory*, Springer - Verlag (1970)
10. J.Roe, *Lectures On Coarse Geometry*, Univ. Lect. Series Vol. 31 Amer.Math.Society.(2003)
11. J.P.Serre, *Trees*, Springer-Verlag (1980)
12. J.Smith, *On Asymptotic Dimension Of Countable Abelian Groups*, Topology and Applications, Vol 153, No 12, 2047, June (2006)
13. G.Yu, *The Novikov conjecture for groups with finite asymptotic dimension*, Ann. of Math. 147(1998), No 2, 325-335.

MATHEMATICS DEPARTMENT, UNIVERSITY OF ATHENS, ATHENS 157 84, GREECE
E-mail address: gentimisth@math.uoa.gr